

Measuring the Hausdorff Dimension of Quantum Mechanical Paths

H. Kröger^a, S. Lantagne^a, K.J.M. Moriarty^b and B. Plache^b

^a Département de Physique, Université Laval,
Québec, Québec G1K 7P4, Canada

^b Department of Mathematics, Statistics and Computing Science
Dalhousie University, Halifax, Nova Scotia B3H 3J5, Canada

Université Laval preprint LAVAL-PHY 7/94
October 1994

Abstract

We measure the propagator length in imaginary time quantum mechanics by Monte Carlo simulation on a lattice and extract the Hausdorff dimension d_H . We find that all local potentials fall into the same universality class giving $d_H = 2$ like the free motion. A velocity dependent action ($S \propto \int dt |\vec{v}|^\alpha$) in the path integral (e.g. electrons moving in solids, or Brueckner's theory of nuclear matter) yields $d_H = \frac{\alpha}{\alpha-1}$ if $\alpha > 2$ and $d_H = 2$ if $\alpha \leq 2$. We discuss the relevance of fractal pathes in solid state physics and in QFT , in particular for the Wilson loop in QCD .

PACS index: 03.65.-w, 05.30.-d

When considering the geometry of particle propagation in quantum mechanics the suitable language is that of path integrals. Feynman and Hibbs [1] have shown that quantum mechanical paths can be viewed as zig-zag lines, which are nowhere differentiable, and which exhibit self-similarity when viewed at different length scales. Abbot and Wise [2] have shown that quantum mechanical paths of free motion are fractal curves [3] of Hausdorff dimension two. In imaginary time (Euclidean) quantum mechanics the free motion resembles very much the Brownian motion of a classical particle as has been observed by Nelson [4]. It turns out that the average path of a classical particle carrying out a Brownian motion (steps Δx and Δt , such that the diffusion coefficient $d = \frac{1}{2}\Delta x^2/\Delta t$ is finite) is also a fractal curve of Hausdorff dimension two [5, 6]. The Euclidean path integral of imaginary time quantum mechanics corresponds to a Wiener measure [5]. The case of the Euclidean free scalar field is discussed by Itzykson and Drouffe [6]. Quite generally, the measure of Euclidean path integrals gives the dominant contributions coming from field configurations corresponding to paths of nowhere differentiable curves [7]. By measuring the length L of a curve in terms of an elementary length Δx , the geometrical property of being a fractal is captured in the Hausdorff dimension d_H , defined by

$$L \sim_{\Delta x \rightarrow 0} L_0 (\Delta x)^{1-d_H}. \quad (1)$$

Strictly speaking, this gives a fractal dimension d_f which in most cases coincides with the Hausdorff dimension d_H . In quantum mechanics, $\Delta x \rightarrow 0$ requires $\Delta t \rightarrow 0$, one is at the continuum limit. The Hausdorff dimension is a critical exponent.

In this letter we want to investigate the critical exponent d_H of quantum mechanics in the presence of local potentials and velocity-dependent potentials in the action of the path integral via Monte Carlo simulations on a temporal lattice. In order to be able to do a Monte Carlo simulation we consider imaginary (Euclidean) quantum mechanics with a negative action

$$S[\vec{x}, \epsilon] = -\frac{1}{\hbar} \sum_{j=0}^{N-1} \epsilon \frac{m}{2} \left(\frac{\vec{x}_{j+1} - \vec{x}_j}{\epsilon} \right)^2, \quad (2)$$

where we have discretized the time $T = t_f - t_i$, $\Delta t = T/N \equiv \epsilon$. We define the quantum mechanical expectation value of the length for propagating the particle from $(\vec{x}_i, t_i)$ to

$(\vec{x}_f, t_f)$ in D space dimensions via

$$\langle L_{f,i}(\epsilon) \rangle = \langle \sum_{k=0}^{N-1} |\vec{x}_{k+1} - \vec{x}_k| \rangle = \frac{\int d^D x_1 \cdots d^D x_{N-1} \sum_{k=0}^{N-1} |\vec{x}_{k+1} - \vec{x}_k| \exp[S[\vec{x}, \epsilon]]}{\int d^D x_1 \cdots d^D x_{N-1} \exp[S[\vec{x}, \epsilon]]} \Big|_{\vec{x}_0=\vec{x}_i, \vec{x}_N=\vec{x}_f} . \quad (3)$$

It is useful to introduce another quantity related to the length squared

$$\langle SQ_{f,i}(\epsilon) \rangle = \langle \sum_{k=0}^{N-1} (\vec{x}_{k+1} - \vec{x}_k)^2 \rangle . \quad (4)$$

This can be computed analytically

$$\langle SQ_{f,i}(\epsilon) \rangle = \frac{D\hbar\epsilon}{2m} \left[\frac{2m(\vec{x}_f - \vec{x}_i)^2}{D\hbar T} - 2 \right] + \frac{D\hbar T}{m} . \quad (5)$$

In the continuum limit ($\Delta t \rightarrow 0$, $|\Delta \vec{x}| \rightarrow 0$), we expect critical behavior

$$\langle L_{f,i} \rangle \sim (\Delta t)^{-A} \sim (\Delta x)^{1-d_H} , \quad (6)$$

$$\langle SQ_{f,i} \rangle \sim (\Delta t)^{-B} \sim (\Delta x)^C . \quad (7)$$

In order to determine the Hausdorff dimension, one needs to know the elementary length Δx . From the action one can estimate this in $D = 1$ dimension from $\frac{\Delta t}{\hbar} \frac{m}{2} (\Delta x / \Delta t)^2 \approx 1$, to give in D dimensions $\langle |\Delta \vec{x}| \rangle_{est} \approx \sqrt{2D\hbar\Delta t/m}$. From that we can estimate $\langle L_{f,i} \rangle$ and $\langle SQ_{f,i} \rangle$

$$\langle L_{f,i}(\epsilon) \rangle_{est} = N \langle |\Delta \vec{x}| \rangle = T \sqrt{2D\hbar/m\Delta t} = 2D\hbar T/m (\langle |\Delta \vec{x}| \rangle)^{-1} , \quad (8)$$

$$\langle SQ_{f,i}(\epsilon) \rangle_{est} = N \langle (\Delta \vec{x})^2 \rangle = 2D\hbar T/m . \quad (9)$$

Comparing eq.(8) with eq.(6) yields $A = 1/2$ and $d_H = 2$ confirming the known result for free motion. Eq.(9) is in agreement (apart from an overall factor 2) with the exact result of eq.(5) (for $\epsilon \rightarrow 0$) and yields $B = C = 0$.

Results of $\langle L_{f,i} \rangle$ and $\langle SQ_{f,i} \rangle$ from Monte Carlo simulation are shown in Fig.[1]. We have considered in $D = 3$ space dimensions the free motion as well as motion in the presence of a local potential $V(\vec{x})$, i.e. the cases: (a) $V = 0$ (free motion), (b) $V = V_0 \vec{x}^2$ (harmonic oscillator), (c) $V = V_0 / |\vec{x}|$ (Coulomb potential). We have plotted $\log \langle L \rangle$

versus $\log N$ and $\langle SQ \rangle$ versus N , where $T = N\Delta t$ is kept fixed. The expectation value $\langle |\Delta \vec{x}| \rangle$ is obtained by $\langle L \rangle / N$. In Fig.[1a,b] we show $\langle L \rangle$ and $\langle SQ \rangle$ for the free case and find agreement with the estimated scaling behavior given by eq.(6-9). In Fig.[1c] we display $\langle L \rangle$ for the harmonic oscillator. Within numerical errors we find for the free motion as well as for both local potentials the same slope. We have extracted the Hausdorff dimension d_H which in all cases is found to be consistent within numerical errors with $d_H = 2$. Thus in the language of critical phenomena these local potentials fall all into the same universality class. For the free motion $\langle SQ \rangle$ is consistent with the exact result from eq.(5). Again the presence of the local potentials does not change the critical exponent C which is compatible with zero.

The above estimate of the Hausdorff dimension hinges upon the presence of the velocity squared term in the action. Thus one expects a different fractal dimension if a term with a different velocity dependence is present in the action of the path integral. Such a velocity dependence occurs in solid state physics for electrons moving in solids [8]. Another example in nuclear physics is Brueckner's theory of nuclear matter [9], using a potential of the form $V(k) = V_0 + V_2 k^2 + V_4 k^4$. One should note, however, that a non-Gaussian momentum dependence in the quantum Hamiltonian does *not* correspond to the same functional form in the action of the path integral. Assuming a polynomial momentum dependence, e.g. p^4 , in the Hamiltonian leads in the action to a term which falls off faster than a Gaussian as a function of $\Delta x / \Delta t^{1/4}$. This can be parametrized by $\exp[-(\Delta x / \Delta t^{1/4})^{A_1} + \dots + (\Delta x / \Delta t^{1/4})^{A_M}]$. If $A = 4/3$, this leads to a velocity dependence $\int dt v^{4/3}$ in the action. All terms in the above parametrization lead to a $\Delta x \sim \Delta t$ behavior different from that of the free motion $\Delta x \sim \sqrt{\Delta t}$. In order to keep matters simple, we have chosen to consider the following parametrization of the action of the path integral

$$S = - \int dt \frac{1}{2} m \vec{v}^2 + V_0 |\vec{v}|^\alpha + U(\vec{x}). \quad (10)$$

Here U denotes a local potential. We have seen above that a local potential does not change the fractal dimension, so we drop the U term in what follows. Let us estimate the Hausdorff dimension for the action $S = - \int dt V(\vec{v})$. In analogy to the free case we have

$\Delta t V_0(<|\Delta\vec{x}|> / \Delta t)^\alpha \propto 1$, and thus

$$<|\Delta\vec{x}|> \propto (\Delta t)^{\frac{\alpha-1}{\alpha}}. \quad (11)$$

This implies

$$<L_{f,i}(\epsilon)>_{est} \propto (\Delta t)^{-\frac{1}{\alpha}} \propto <|\Delta\vec{x}|>_{est}^{-\frac{1}{\alpha-1}}, \quad (12)$$

$$<SQ_{f,i}(\epsilon)>_{est} \propto (\Delta t)^{\frac{\alpha-2}{\alpha}} \propto <|\Delta\vec{x}|>_{est}^{\frac{\alpha-2}{\alpha-1}}, \quad (13)$$

Of course, for $\alpha = 2$ the scaling behavior agrees with that of the free motion, eqs.(8,9). From eq.(12), we imply for the Hausdorff dimension $d_H = \frac{\alpha}{\alpha-1}$ and for the critical exponent $C = \frac{\alpha-2}{\alpha-1}$. This opens the mathematical possibility of having a very large Hausdorff dimension, if α is above but close to unity. However, physically this is not realized, because the presence of the kinetic term in the action limits $d_H \leq 2$, as will be shown below.

The results of the Monte Carlo simulations with action given by eq.(10) corresponding to $D = 3$ dimensions are shown in Figs.[2,3] In Fig.[2] we display $<L>$ and $<SQ>$ versus $\log N$ for $\alpha = 1/2$. We have extracted the critical exponents d_H and C as a function of α and show a plot in Fig.[3]. We find within numerical errors the following behavior: For $\alpha < 2$, d_H agrees with the value of the free motion, i.e., $d_H = 2$, while for $\alpha > 2$, d_H agrees with the value $\frac{\alpha}{\alpha-1}$, i.e., the value expected from the velocity dependent potential without kinetic energy. Thus $\alpha_{crit} = 2$ is a critical value of the velocity dependent potential. The same kind of behavior is seen for the critical exponent C .

To conclude let us discuss some perspectives for solid state physics and for *QCD*. Consider in solid state physics a semiconductor where conduction properties (due to doping) change with position, e.g., like in a transistor. The Hausdorff dimension is sensitive to higher order terms in the energy-momentum dispersion relation. Although the determination of d_H needs a non-local observable, i.e., $<L>$, the absolute length is not relevant for d_H . So in principle one could measure $<L>$ on a length scale large to atomic distance, but small to the length of variation of macroscopic properties. In such a way, one could determine d_H locally and study its change as a function of position throughout the semiconductor (this would correspond to a velocity dependent potential, where α is a function of position).

In this sense d_H can be viewed as one parameter probing the local geometry of quantum mechanics. In QCD , a fractal dimension might eventually play a role in a non-local observable, like the Wilson-loop. The Wilson loop distinguishes between a confined (area law) and a non-confined (perimeter law) phase. It also has a quark-quark potential interpretation which yields the string tension, being crucial for the baryon spectrum. The potential interpretation makes the semiclassical assumption that the quarks propagate along classical smooth curves. In quantum mechanics however, one would expect the quarks to propagate along zig-zag lines. The path of quark propagation enters in the definition of the Wilson loop. Going over from smooth curves, $d_H = 1$, to zig-zag lines, e.g., with $d_H = 2$ which are similar to a surface, would most likely change the behavior of this observable.

Acknowledgement H.K. is grateful to J. Polonyi and V. Branchina for helpful discussions. H.K. was supported by NSERC Canada and FCAR Québec.

References

- [1] R.P. Feynman and A.R. Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill, New York (1965).
- [2] L.F. Abbot and M.B. Wise, Am. J. Phys. 49 (1981) 37.
- [3] B.B. Mandelbrot, The Fractal Geometry of Nature, Freeman, New York (1983).
- [4] E. Nelson, Phys. Rev. 150 (1966) 1079.
- [5] L.S. Schulman, Techniques and Applications of Path Integration, Wiley, New York (1981).
- [6] C. Itzykson and J.B. Drouffe, Statistical Field Theory, Cambridge Univ. Press., Cambridge (1989), Vol.I.
- [7] J. Glimm and A. Jaffe, Quantum Physics, Springer, New York (1981).
- [8] O. Madelung, Solid State Theory, Springer, New York (1981).
- [9] K.A. Brueckner, Phys. Rev. 97 (1955) 1353.

Figure Caption


Fig.1 (a) Length $\langle L \rangle$, and (b) sum of length squares $\langle SQ \rangle$ for free motion and (c) length $\langle L \rangle$ for harmonic oscillator versus number of time steps N .


Fig.2 Same as Fig.[1] for velocity dependent action ($\alpha = 1/2$).


Fig.3 Critical exponents d_H and C versus potential parameter α for velocity dependent action. The full line gives the estimated exponents from eq.(11-13).

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>









This figure "fig1-2.png" is available in "png" format from:


<http://arXiv.org/ps/hep-lat/9501018v1>






This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>





This figure "fig1-4.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>

This figure "fig1-5.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>

This figure "fig1-6.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>

This figure "fig1-7.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9501018v1>